



Comparison of Cardiomyogenic Potential of Amniotic Fluid Mesenchymal Stromal Cells Derived from Normal and Isolated Congenital Heart Defective Fetuses

Name: Manali Jain

Affiliation: PhD Scholar, Sanjay Gandhi Post-Graduate Institute of Medical Sciences, Lucknow

Country: India

Email ID: manalijain.sgpgi@gmail.com

ABSTRACT

Isolated Congenital Heart Defect (ICHD) is the major cause of neonatal death worldwide among all forms of CHDs. Recently, stem cell therapies have emerged as a potential approach to ameliorate ICHD in children. ICHD is characterized by cardiac structural abnormalities during embryogenesis due to alterations in a pool of cardiac progenitors/ stem cells associated with fetal heart development. The stem cells present in the amniotic fluid (AF) are of fetal origin and may reflect the physiological and pathological changes in the fetus during embryogenesis. Therefore, in the present study, the cardiomyogenic potential of AF-MSCs derived from fetuses with ICHD (ICHD AF-MSCs) has been evaluated and compared with that of AF-MSCs of structurally normal fetuses (normal AF-MSCs). The immunophenotypic characterization revealed that ICHD AF-MSCs have significantly lower expression of cardiac progenitor markers VEGFR-2 ($p < 0.01$), SSEA-1 ($p < 0.01$), and PDGFR- α ($p < 0.01$) in comparison to normal AF-MSCs. Upon induction with 5'-azacytidine for 21 days, ICHD AF-MSCs showed a significantly down-regulated expression of

cardiac transcription factors such as GATA-4 (0.4 ± 0.1 vs. 6.8 ± 1.2 ; $p < 0.01$), ISL-1 (2.3 ± 0.6 vs. 14.3 ± 1.12 ; $p < 0.01$), NK-x 2-5 (1.1 ± 0.3 vs. 14.1 ± 2.8 ; $p < 0.01$), TBX-5 (0.4 ± 0.07 vs. 4.4 ± 0.3 ; $p < 0.001$), and TBX-18 (1.3 ± 0.2 vs. 4.19 ± 0.3 ; $p < 0.01$) when compared with the normal AF-MSCs. Furthermore, immunocytochemical staining revealed that both types of AF-MSCs could differentiate into cardiovascular lineages and express cardiomyogenic, endothelial, and smooth muscle actin markers, viz., cardiac troponin (cTNT), CD31, and alpha-smooth muscle actin (α -SMA). However, normal AF-MSCs showed an enhanced expression of cTNT ($p < 0.001$), CD31 ($p < 0.01$), and α -SMA ($p < 0.05$), compared to ICHD AF-MSCs. Overall, these results suggest that the ICHD-AF-MSCs have a defective cardiomyogenic differentiation potential and that the defects in these stem cells may have a role in the pathogenesis of ICHD.

BIOGRAPHY

Manali Jain is a PhD Scholar in the department of Hematology, Sanjay Gandhi Post-Graduate Institute of Medical Sciences, Lucknow, India. She is presently working in the field of cardiac regenerative potential of amniotic fluid stem cells.